

UCAS PERSONAL ID:

NAME:

OXFORD UNIVERSITY

MATHEMATICS ADMISSIONS TEST

Wednesday 2 November 2011

Time Allowed: 21/2 hours

For candidates applying for Mathematics, Computer Science or one of their joint degrees.

Write your name, test centre (where you are sitting the test), Oxford college (to which you applied or were assigned) and your proposed degree course in BLOCK CAPITALS below.


OXFORD COLLEGE (if know	wn):								
DEGREE COURSE									
DATE OF BIRTH:									
This paper contains 7 question paper as to which questions a	ns of which are appropri	you sho iate for yo	uld attem our cours	pt 5. The	re are dire	ctions th	roughout the		
Mathematics, Maths & Mathematics & Compu Computer Science, Co	ter Science mputer Scie	applican ence & Ph	ts should iilosophy	attempt applican	1,2,3,5,6.				
Further credit cannot be obtained by attempting extra questions. Question 1 is a multiple choice question with ten parts. Marks are given solely for correct answers but any rough working should be shown in the space between parts. Answer Question 1 on the grid on Page 2. Each part is worth 4 marks.									
Answers to questions 2-7 should be written in the space provided, continuing on to the blank pages at the end of this booklet if necessary. Each of Questions 2-7 is worth 15 marks.									
For Test Supervisors Use Onl	V*								
[] Tick here if special arrang securely attach to the test scr	gements we	ere made with the d	for the te letails.	st. Please	either ind	clude deta	ails below or		
Signature of Invigilator:									
FOR OFFICE USE ONLY:	Q1	Q2	Q3	Q4	Q5	Q6	Q7		
		· ·							


1. For ALL APPLICANTS.

For each part of the question on pages 3–7 you will be given four possible answers, just one of which is correct. Indicate for each part A-J which answer (a), (b), (c), or (d) you think is correct with a tick (\checkmark) in the corresponding column in the table below. Please show any rough working in the space provided between the parts.

	(a)	(b)	(c)	(d)
A				
В				
C		×		2
D				
E	,			
F	¥i			
G				
Н				
I				
J				

A. A sketch of the graph $y = x^3 - x^2 - x + 1$ appears on which of the following axes?





B. A rectangle has perimeter P and area A. The values P and A must satisfy

- (a) $P^3 > A$, (b) $A^2 > 2P + 1$, (c) $P^2 \ge 16A$, (d) $PA \ge A + P$.

Turn over

C. The sequence x_n is given by the formula

$$x_n = n^3 - 9n^2 + 631.$$


- The largest value of n for which $x_n > x_{n+1}$ is
 - (a) 5,
- (b) 7, (c) 11,
- (d) 17.

D. The fraction of the interval $0 \le x \le 2\pi$, for which one (or both) of the inequalities

$$\sin x \geqslant \frac{1}{2}, \qquad \sin 2x \geqslant \frac{1}{2}$$

- is true, equals
- (a) $\frac{1}{3}$, (b) $\frac{13}{24}$, (c) $\frac{7}{12}$, (d) $\frac{5}{8}$.

E. The circle in the diagram has centre C. Three angles α, β, γ are also indicated.



The angles α, β, γ are related by the equation:

- (a) $\cos \alpha = \sin (\beta + \gamma)$;
- (b) $\sin \beta = \sin \alpha \sin \gamma$;
- (c) $\sin \beta (1 \cos \alpha) = \sin \gamma;$
- (d) $\sin(\alpha + \beta) = \cos \gamma \sin \alpha$.
- F. Given θ in the range $0 \le \theta < \pi$, the equation


$$x^2 + y^2 + 4x \cos \theta + 8y \sin \theta + 10 = 0$$

represents a circle for

- (a) $0 < \theta < \frac{\pi}{3}$, (b) $\frac{\pi}{4} < \theta < \frac{3\pi}{4}$, (c) $0 < \theta < \frac{\pi}{2}$, (d) all values of θ .

Turn over

G. A graph of the function y = f(x) is sketched on the axes below:



- The value of $\int_{-1}^{1} f(x^2 1) dx$ equals

- (a) $\frac{1}{4}$, (b) $\frac{1}{3}$, (c) $\frac{3}{5}$, (d) $\frac{2}{3}$.

 \mathbf{H} . The number of positive values x which satisfy the equation

$$x = 8^{\log_2 x} - 9^{\log_3 x} - 4^{\log_2 x} + \log_{0.5} 0.25$$

is

- (a) 0,
- (b) 1,
- (c) 2,

I. In the range $0 \leqslant x < 2\pi$ the equation

$$\sin^8 x + \cos^6 x = 1$$

has

- (a) 3 solutions,
- (b) 4 solutions,
- (c) 6 solutions,
- (d) 8 solutions.

J. The function f(n) is defined for positive integers n according to the rules

$$f(1) = 1,$$

$$f(2n) = f(n)$$

$$f(2n) = f(n),$$
 $f(2n+1) = (f(n))^2 - 2.$

The value of $f(1) + f(2) + f(3) + \cdots + f(100)$ is

(a)
$$-86$$
, (b) -31 , (c) 23 ,

(b)
$$-31$$

Turn over

2. For ALL APPLICANTS.

Suppose that x satisfies the equation

$$x^3 = 2x + 1. (*)$$

(i) Show that

$$x^4 = x + 2x^2$$
 and $x^5 = 2 + 4x + x^2$.

(ii) For every integer $k \geqslant 0$, we can uniquely write

$$x^k = A_k + B_k x + C_k x^2$$

where A_k , B_k , C_k are integers. So, in part (i), it was shown that

$$A_4 = 0, B_4 = 1, C_4 = 2$$
 and $A_5 = 2, B_5 = 4, C_5 = 1$.

Show that

$$A_{k+1} = C_k$$
, $B_{k+1} = A_k + 2C_k$, $C_{k+1} = B_k$.

(iii) Let


$$D_k = A_k + C_k - B_k.$$

Show that $D_{k+1} = -D_k$ and hence that

$$A_k + C_k = B_k + (-1)^k$$
.

(iv) Let $F_k = A_{k+1} + C_{k+1}$. Show that

$$F_k + F_{k+1} = F_{k+2}$$
.



For APPLICANTS IN


MATHEMATICS
MATHEMATICS & STATISTICS
MATHEMATICS & PHILOSOPHY
MATHEMATICS & COMPUTER SCIENCE

ONLY.

Computer Science and Computer Science & Philosophy applicants should turn to page 14.

The graphs of $y = x^3 - x$ and y = m(x - a) are drawn on the axes below. Here m > 0 and $a \le -1$.

The line y = m(x - a) meets the x-axis at A = (a, 0), touches the cubic $y = x^3 - x$ at B and intersects again with the cubic at C. The x-coordinates of B and C are respectively b and c.



- (i) Use the fact that the line and cubic touch when x = b, to show that $m = 3b^2 1$.
- (ii) Show further that

$$a = \frac{2b^3}{3b^2 - 1}.$$


- (iii) If $a = -10^6$, what is the approximate value of b?
- (iv) Using the fact that

$$x^{3} - x - m(x - a) = (x - b)^{2}(x - c)$$

(which you need not prove), show that c = -2b.

(v) R is the finite region bounded above by the line y = m(x - a) and bounded below by the cubic $y = x^3 - x$. For what value of a is the area of R largest?

Show that the largest possible area of R is $\frac{27}{4}$.



4.


For APPLICANTS IN

MATHEMATICS MATHEMATICS & STATISTICS MATHEMATICS & PHILOSOPHY

ONLY.

Mathematics & Computer Science, Computer Science and Computer Science & Philosophy applicants should turn to page 14.

Let Q denote the quarter-disc of points (x,y) such that $x \ge 0$, $y \ge 0$ and $x^2 + y^2 \le 1$ as drawn in Figures A and B below.



(i) On the axes in Figure A, sketch the graphs of

$$x + y = \frac{1}{2}$$
, $x + y = 1$, $x + y = \frac{3}{2}$.

What is the largest value of x + y achieved at points (x, y) in Q? Justify your answer.

(ii) On the axes in Figure B, sketch the graphs of

$$xy = \frac{1}{4}$$
, $xy = 1$, $xy = 2$.

What is the largest value of $x^2 + y^2 + 4xy$ achieved at points (x, y) in Q?

What is the largest value of $x^2 + y^2 - 6xy$ achieved at points (x, y) in Q?

(iii) Describe the curve

$$x^2 + y^2 - 4x - 2y = k$$

where k > -5.

What is the smallest value of $x^2 + y^2 - 4x - 2y$ achieved at points (x, y) in Q?



5. For ALL APPLICANTS.

An $n \times n$ grid consists of squares arranged in n rows and n columns; for example, a chessboard is an 8×8 grid. Let us call a semi-grid of size n the lower left part of an $n \times n$ grid – that is, the squares located on or below the grid's diagonal. For example, Figure C shows an example of a semi-grid of size 4.




Figure C

Let us suppose that a robot is located in the lower-left corner of the grid. The robot can move only up or right, and its goal is to reach one of the goal squares, which are all located on the semi-grid's diagonal. In the example shown in Figure C, the robot is initially located in the square denoted with R, and the goal squares are shown in grey. Let us call a solution a sequence of the robot's moves that leads the robot from the

- (i) Write down all 8 solutions for a robot on a semi-grid of size 4.
- (ii) Devise a concise way of representing the possible journeys of the robot in a semi-grid of size n. In your notation, which of the journeys are solutions?
- (iii) Write down a formula for the number of possible solutions in a semi-grid of size n.

Now let us change the problem slightly and redefine a goal square as any square that

- the lower-left square is not a goal square;
- each square that is located immediately above or immediately to the right of a non-goal square is a goal square; and
- each square that is located immediately above or immediately to the right of a goal square is a non-goal square.

Furthermore, let us assume that, upon reaching a goal square, the robot may decide to stop or to continue moving (provided that there are more allowed moves).

- (iv) With these modifications in place, write down all the solutions in a semi-grid of size 4, and all the solutions in a semi-grid of size 5.
- (v) How many solutions are there now in a semi-grid of size n, where n is a positive integer? You may wish to consider separately the cases where n is even or odd.



6.

For APPLICANTS IN COMPUTER SCIENCE MATHEMATICS & COMPUTER SCIENCE COMPUTER SCIENCE & PHILOSOPHY ONLY.

Alice, Bob, Charlie and Diane are playing together when one of them breaks a precious vase. They all know who broke the vase. When questioned they make the following statements:

Alice: It was Bob.
Bob: It was Diane.
Charlie: It was not me.

Diane: What Bob says is wrong.

Each statement is either true or false.

- (i) Explain why at least one of the four must be lying.
- (ii) Explain why at least one of them must be telling the truth.
- (iii) Let us suppose that exactly one of the four is lying, so the other three are telling the truth. Who is lying? Who did break the vase? Explain your answer.
- (iv) Let us now suppose that exactly one of the four is telling the truth, so the other three are lying. Who is telling the truth? Who did break the vase? Explain your answer.
- (v) Let us now suppose that two of the statements are true and two are false. List the people who might now have broken the vase. Justify your answers.
- (vi) Hence show that if we don't know how many of the four statements are true, then any one of the four could have broken the vase.

Turn over

7. For APPLICANTS IN $\left\{ \begin{array}{l} \text{COMPUTER SCIENCE} \\ \text{COMPUTER SCIENCE \& PHILOSOPHY} \end{array} \right\}$ ONLY.

Alice and Bob have a large bag of coins which they use to play a game called HT-2. In this game, Alice and Bob take turns placing one coin at a time on the table, each to the right of the previous one; thus they build a row of coins that grows to the right. Alice always places the first coin. Each coin is placed head-up (H) or tail-up (T), and cannot be flipped or moved once it has been placed.

A player loses the game if he or she places a coin that results in two adjacent coins having the same pattern of heads and tails as another adjacent pair somewhere in the row (reading a pattern from left to right). For example, Bob has lost this game by producing a second instance of HT (where a and b denote coins placed by Alice and Bob respectively):

and Alice has lost this game by producing a second instance of TT (overlapping pairs can count as repeats):

- (i) What is the smallest number of coins that might be placed in a game of HT-2 (including the final coin that causes a player to lose)? What is the largest number? Justify each answer.
- (ii) Bob can always win a game of HT-2 by adopting a particular strategy. Describe the strategy.

For any positive integer n, there is a game HT-n with the same rules as HT-2, except that the game is lost by a player who creates an unbroken sequence of n heads and tails that appears elsewhere in the row. For example, Bob has lost this game of HT-3 by producing a second instance of THT:

(iii) Suppose n is odd, and Bob chooses to play by always duplicating Alice's previous play (and Alice knows that this is Bob's strategy). Show that Alice can always win.

In these games, a maximum time of one minute is allowed for each turn.

(iv) Can we be certain that a game of HT-6 will be finished within two hours? Justify your answer.

End of last question

Latin Unseen Translation

Time allowed: 1 hour

Translate both passages into English. Use the answer spaces provided under each question.

Special provisions (e.g. extra time)

UCAS Personal ID

Surname

First name(s)

Date of birth

 A farmer entertains in his cottage three gods travelling incognito, until Neptune inadvertently gives them away.

luppiter et lato qui regnat in aequore frater carpebant socias Mercuriusque vias;

forte senex Hyrieus, angusti cultor agelli,

hos videt, exiguam stabat ut ante casam,

atque ita 'longa via est, nec tempora longa supersunt',

dixit 'et hospitibus ianua nostra patet.'

tecta senis subeunt nigro deformia fumo;

ignis in hesterno stipite parvus erat.

dumque mora est, tremula dat vina rubentia dextra:

accipit aequoreus pocula prima deus.

quae simul exhausit, 'da nunc bibat ordine' dixit

'luppiter.' audito palluit ille love.

Ovid Fasti 5. 495-514 (selected verses)

carpo (+ acc.) hurry along

stipes log

2. A gigantic Gaul challenges the Romans to produce a warrior for single combat.

Tum Gallus eximia corporis magnitudine processit et maxima voce 'quem nunc' inquit 'Roma virum fortissimum habet, procedat ad pugnam, ut eventus ostendat utra gens sit bello melior.' Iuvenem armatum Romani adversus Gallum producunt. Recipiunt inde se ad stationem; et duo in medio relinquuntur, nequaquam pares. Corpus alteri magnitudine erat eximium, versicolori veste pictisque refulgens armis; media alterius statura armaque habilia magis quam decora. Ubi constitere inter duas acies, Gallus in advenientis arma hostis cum ingenti sonitu ensem deiecit.

Livy 7.9 (adapted)

nequaquam in no respect

versicolor multicoloured

habilis serviceable

Then the Gawl when they steed a more serviced in the man laborate and how he man as its the strongest, let him come out end to fight, so that the sent in war. "The lamans led and amed your against the Gawl. From there they welcomed him into his place; and they left the two of one was greatly enominated armour; the other was hard his height and his weapons were more serviceable than elaborate, when they steed there was hard elaborate, when they steed there was between the power in the property when elaborate when they steed there was between the power in they steed there weapons of the enomy as he was approaching, the Gawl threw enomy as he was approaching, the Gawl threw